
Discrete Structures

CSE 2315 (Spring 2014)

Lecture 17 Graphs

Heng Huang, Ph.D.

Department of Computer Science and Engineering

Agenda – Graphs

- Graph basics and definitions
 - Vertices/nodes, edges, adjacency, incidence
 - Degree, in-degree, out-degree
 - Degree, in-degree, out-degree
 - Subgraphs, unions, isomorphism
 - Adjacency matrices
- Types of Graphs
 - Trees
 - Undirected graphs
 - Simple graphs, Multigraphs, Pseudographs
 - Digraphs, Directed multigraph
 - Bipartite
 - Complete graphs, cycles, wheels, cubes, complete bipartite

Uses of Graph Theory in CS

- Car navigation system
- Efficient database
- Build a bot to retrieve info off WWW
- Representing computational models
- Many other applications.
- This course we focus more on the properties of abstract graphs rather on algorithms

Graphs –Intuitive Notion

A graph is a bunch of vertices (or nodes) represented by circles • which are connected by edges, represented by line segments .

Mathematically, graphs are binary-relations on their vertex set (except for multigraphs).

In Data Structures one often starts with trees and generalizes to graphs. In this course, opposite approach: We start with graphs and restrict to get trees.


Trees

A very important type of graph in CS is called a *tree*:




Trees

A very important type of graph in CS is called a *tree*:




Trees

A very important type of graph in CS is called a *tree*:



Trees

A very important type of graph in CS is called a *tree*:



Real
Tree

Abstract
Tree

Simple Graphs


Different purposes require different types of graphs.

EG: Suppose a local computer network

- Is bidirectional (undirected)
- Has no loops (no “self-communication”)
- Has unique connections between computers

Sensible to represent as follows:

Simple Graphs



- Vertices are labeled to associate with particular computers
- Each edge can be viewed as the set of its two endpoints

Simple Graphs

DEF: A *simple graph* $G = (V, E)$ consists of a non-empty set V of *vertices* (or *nodes*) and a set E (possibly empty) of *edges* where each edge is a subset of V with cardinality 2 (an unordered pair).

Q: For a set V with n elements, how many possible edges there?

Simple Graphs

A: The number of pairs in V

$$= C(n, 2) = n \cdot (n - 1) / 2$$

Q: How many possible graphs are there for the same set of vertices V ?


Simple Graphs

A: The number of subsets in the set of possible edges.
There are $n \cdot (n - 1) / 2$ possible edges, therefore the number of graphs on V is $2^{n(n-1)/2}$

Multigraphs

If computers are connected via internet instead of directly, there may be several routes to choose from for each connection. Depending on traffic, one route could be better than another. Makes sense to allow multiple edges, but still no self-loops:

Multigraphs



Edge-labels distinguish between edges sharing same endpoints. Labeling can be thought of as function:


$$\begin{aligned} e_1 &\rightarrow \{1,2\}, e_2 \rightarrow \{1,2\}, e_3 \rightarrow \{1,3\}, \\ e_4 &\rightarrow \{2,3\}, e_5 \rightarrow \{2,3\}, e_6 \rightarrow \{1,2\} \end{aligned}$$

Multigraphs

DEF: A *multigraph* $G = (V, E, f)$ consists of a non-empty set V of *vertices* (or *nodes*), a set E (possibly empty) of *edges* and a function f with domain E and codomain the set of pairs in V .

Pseudographs

If self-loops are allowed we get a pseudograph:



Now edges may be associated with a single vertex, when the edge is a loop

$$e_1 \rightarrow \{1,2\}, e_2 \rightarrow \{1,2\}, e_3 \rightarrow \{1,3\},$$


$$e_4 \rightarrow \{2,3\}, e_5 \rightarrow \{2\}, e_6 \rightarrow \{2\}, e_7 \rightarrow \{4\}$$

Multigraphs

DEF: A *pseudograph* $G = (V, E, f)$ consists of a non-empty set V of *vertices* (or *nodes*), a set E (possibly empty) of *edges* and a function f with domain E and codomain the set of pairs and singletons in V .


Undirected Graphs Terminology

Vertices are *adjacent* if they are the endpoints of the same edge.



Q: Which vertices are adjacent to 1? How about adjacent to 2, 3, and 4?

Undirected Graphs Terminology



A: 1 is adjacent to 2 and 3


2 is adjacent to 1 and 3

3 is adjacent to 1 and 2

4 is not adjacent to any vertex


Undirected Graphs Terminology

A vertex is *incident* with an edge (and the edge is incident with the vertex) if it is the endpoint of the edge.



Q: Which edges are incident to 1? How about incident to 2, 3, and 4?

Undirected Graphs Terminology



A: e_1, e_2, e_3, e_6 are incident with 1


2 is incident with e_1, e_2, e_4, e_5, e_6

3 is incident with e_3, e_4, e_5

4 is not incident with any edge

Digraphs


Digraphs can be used as a way of representing relations:



Q: What type of pair should each edge be (multiple edges not allowed)?

Digraphs

A: Each edge is directed so an *ordered* pair (or tuple) rather than unordered pair.



Thus the set of edges E is just the represented relation on V .

Digraphs

DEF: A *directed graph* (or *digraph*) G
= (V, E) consists of a non-empty set V of *vertices*
(or *nodes*) and a set E of *edges* with $E \subseteq V \times V$.

The edge (a, b) is also denoted by $a \rightarrow b$ and a is called the *source* of the edge while b is called the *target* of the edge.


Q: For a set V with n elements, how many possible digraphs are there?

Digraphs

A: The same as the number of relations on V , which is the number of subsets of $V \times V$ so $2^{n \cdot n}$.

Directed Multigraphs


If also want to allow multiple edges in a digraph, get a *directed multigraph* (or *multi-digraph*).



Q: How to use sets and functions to deal with multiple directed edges, loops?

Directed Multigraphs


A: Have function with domain the edge set and codomain $V \times V$.



$e_1 \rightarrow (1,2), e_2 \rightarrow (1,2), e_3 \rightarrow (2,2), e_4 \rightarrow (2,3),$
 $e_5 \rightarrow (2,3), e_6 \rightarrow (3,3), e_7 \rightarrow (3,3)$


Degree

The ***degree*** of a vertex counts the number of edges that *seem* to be sticking out if you looked under a magnifying glass:




Degree

The ***degree*** of a vertex counts the number of edges that *seem* to be sticking out if you looked under a magnifying glass:



Degree

The ***degree*** of a vertex counts the number of edges that *seem* to be sticking out if you looked under a magnifying glass:




Thus $\deg(2) = 7$ even though 2 only incident with 5 edges.

Q: How to define this formally?


Degree

A: Add 1 for every regular edge incident with vertex and 2 for every loop. Thus $\deg(2) = 1 + 1 + 1 + 2 + 2 = 7$



Oriented Degree when Edges Directed

The ***in-degree*** of a vertex ($\deg^-$) counts the number of edges that stick *in* to the vertex. The ***out-degree*** ($\deg^+$) counts the number sticking *out*.




Q: What are in-degrees and out-degrees of all the vertices?


Oriented Degree when Edges Directed

A:

$$\begin{aligned}\deg^-(1) &= 0 \\ \deg^-(2) &= 3 \\ \deg^-(3) &= 4 \\ \deg^+(1) &= 2 \\ \deg^+(2) &= 3 \\ \deg^+(3) &= 2\end{aligned}$$



Handshaking Theorem



There are two ways to count the number of edges in the above graph:

1. Just count the set of edges: 7
2. Count *seeming* edges vertex by vertex and divide by 2 because double-counted edges:

$$(\deg(1) + \deg(2) + \deg(3) + \deg(4)) / 2 = (3 + 7 + 2 + 2) / 2 = 14 / 2 = 7$$

Handshaking Theorem

THM: In an undirected graph

$$|E| = \frac{1}{2} \sum_{e \in E} \deg(e)$$

In a directed graph

$$|E| = \sum_{e \in E} \deg^+(e) = \sum_{e \in E} \deg^-(e)$$

Q: In a party of 5 people can each person be friends with exactly three others?

Handshaking Theorem

A: Imagine a simple graph with 5 people as *vertices* and edges being undirected edges between friends (simple graph assuming friendship is symmetric and irreflexive). Number of friends each person has is the degree of the person.

Handshaking would imply that

$$|E| = (\text{sum of degrees})/2 \quad \text{or}$$

$$2|E| = (\text{sum of degrees}) = (5 \cdot 3) = 15.$$

Impossible as 15 is not even. In general:

Handshaking Theorem

Lemma: The number of vertices of odd degree must be even in an undirected graph.

Proof: Otherwise would have

$$2|E| = \text{Sum of even no.'s} \\ + \text{an odd number of odd no.'s}$$

$$\Rightarrow \text{even} = \text{even} + \text{odd}$$


—this is impossible. •

Graph Patterns Complete Graphs - K_n


A simple graph is ***complete*** if every pair of distinct vertices share an edge. The notation K_n denotes the complete graph on n vertices.




K_1




K_2



K_3




K_4



K_5

Graph Patterns Cycles - C_n


The *cycle graph* C_n is a circular graph with $V = \{0, 1, 2, \dots, n-1\}$ where vertex i is connected to $i+1 \bmod n$ and to $i-1 \bmod n$. They look like polygons:



Graph Patterns Wheels - W_n

A: Pseudographs


The **wheel graph** W_n is just a cycle graph with an extra vertex in the middle:



Usually consider wheels with 3 or more spokes only.

Graph Patterns Cubes - Q_n

The n -**cube** Q_n is defined recursively. Q_0 is just a vertex.
 Q_{n+1} is gotten by taking 2 copies of Q_n and joining each vertex v of Q_n with its copy v' :




Bipartite Graphs


A simple graph is ***bipartite*** if V can be partitioned into $V = V_1 \cup V_2$ so that any two adjacent vertices are in different parts of the partition. Another way of expressing the same idea is ***bichromatic***: vertices can be colored using two colors so that no two vertices of the same color are adjacent.

Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:




And so is bipartite, if we redraw it:




Bipartite Graphs

EG: C_4 is a bichromatic:



And so is bipartite, if we redraw it:



Q: For which n is C_n bipartite?


Bipartite Graphs

A: C_n is bipartite when n is even. For even n color all odd numbers red and all even numbers green so that vertices are only adjacent to opposite color.


If n is odd, C_n is not bipartite. If it were, color 0 red. So 1 must be green, and 2 must be red. This way, all even numbers must be red, including vertex $n-1$. But $n-1$ connects to 0 $\rightarrow\leftarrow$.

Graph Patterns Complete Bipartite - $K_{m,n}$

When all possible edges exist in a simple bipartite graph with m red vertices and n green vertices, the graph is called ***complete bipartite*** and the notation $K_{m,n}$ is used. EG:




$K_{2,3}$



$K_{4,5}$


Subgraphs

Notice that the 2-cube



occurs

inside the 3-cube



. In other


words, Q_2 is a subgraph of Q_3 :

DEF: Let $G = (V, E)$ and $H = (W, F)$ be graphs. H is said to be a **subgraph** of G , if $W \subseteq V$ and $F \subseteq E$.

Q: How many Q_2 subgraphs does Q_3 have?


Subgraphs

A: Each face of Q_3 is a Q_2 subgraph so the answer is 6, as this is the number of faces on a 3-cube:




Unions

In previous example can actually reconstruct the 3-cube from its 6 2-cube faces:



Unions

If we assign the 2-cube faces (aka S quares) the names $S_1, S_2, S_3, S_4, S_5, S_6$ then Q_3 is the union of its faces:




Unions

DEF: Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two simple graphs (and V_1, V_2 may or may not be disjoint). The ***union*** of G_1, G_2 is formed by taking the union of the vertices and edges. I.E: $G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$.

A similar definitions can be created for unions of digraphs, multigraphs, pseudographs, etc.

Adjacency Matrix

We already saw a way of representing relations on a set with a Boolean matrix:



Adjacency Matrix

Since digraphs *are* relations on their vertex sets, can adopt the concept to represent digraphs. In the context of graphs, we call the representation an ***adjacency matrix***:

For a digraph $G = (V, E)$ define matrix A_G by:

- Rows, Columns –one for each vertex in V
- Value at i^{th} row and j^{th} column is
 - 1 if i^{th} vertex connects to j^{th} vertex ($i \rightarrow j$)
 - 0 otherwise

Adjacency Matrix -Directed Multigraphs


Can easily generalize to directed multigraphs by putting in the number of edges between vertices, instead of only allowing 0 and 1:

For a directed multigraph $G = (V, E)$ define the matrix A_G by:


- Rows, Columns –one for each vertex in V
- Value at i^{th} row and j^{th} column is
 - The number of edges with source the i^{th} vertex and target the j^{th} vertex

Adjacency Matrix -Directed Multigraphs

Q: What is the adjacency matrix?



Adjacency Matrix - Directed Multigraphs




A:

$$\begin{pmatrix} 0 & 3 & 0 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$


Adjacency Matrix -General

Undirected graphs can be viewed as directed graphs by turning each undirected edge into two oppositely oriented directed edges, *except when the edge is a self-loop in which case only 1 directed edge is introduced*. EG:




Adjacency Matrix -General

Q: What's the adjacency matrix?



Adjacency Matrix -General



A:

$$\begin{pmatrix} 0 & 2 & 1 & 0 \\ 2 & 2 & 1 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Notice that answer is *symmetric*.

Adjacency Matrix -General

For an undirected graph $G = (V, E)$ define the matrix A_G by:

- Rows, Columns –one for each element of V
- Value at i^{th} row and j^{th} column is the number of edges incident with vertices i and j .

This is equivalent to converting first to a directed graph as above. Or by allowing undirected edges to take us from i to j can simply use definition for directed graphs.