


Symbol	Notation	Asymptotic Bound	Limit theorem	Definition with constants	Example
Θ ($=$) Theta	$f(n) = \Theta(g(n))$	Asymptotic tight bound ($g(n)$ is an asymptotic tight bound for $f(n)$)	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c \neq 0$ (non-zero constant)	There exist <u>positive</u> constants c_0, c_1 and n_0 s.t.: $c_0 g(n) \leq f(n) \leq c_1 g(n)$ for all $n \geq n_0$	$25n^2 + 100n = \Theta(n^2)$ 
O (\leq) Big-Oh	$f(n) = O(g(n))$	Asymptotic upper bound (can be tight)	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$ or c	There exist <u>positive</u> constants c_1 and n_0 such that: $f(n) \leq c_1 g(n)$ for all $n \geq n_0$	$n^2 + 100n = O(n^3)$ $25n^2 + 100n = O(n^2)$
Ω (\geq) Omega	$f(n) = \Omega(g(n))$	Asymptotic lower bound (can be tight)	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$ or c	There exist <u>positive</u> constants c_0 and n_0 such that: $c_0 g(n) \leq f(n)$ for all $n \geq n_0$	$n^2 + 100n = \Omega(n\sqrt{n})$ $25n^2 + 100n = \Omega(n^2)$ $\frac{n^2}{1000} - 300n = \Omega(n^2)$
o ($<$) Little-oh	$f(n) = o(g(n))$	Asymptotic upper bound but NOT tight	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$ Cannot be a constant	For any <u>positive</u> constant c_1 , there exists n_0 s.t.: $f(n) < c_1 g(n)$ for all $n \geq n_0$	$n^2 + 100n = o(n^3)$ $25n^2 + 100n \neq o(n^2)$
ω ($>$) Little-omega	$f(n) = \omega(g(n))$	Asymptotic lower bound but NOT tight	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$ Cannot be a constant	For any <u>positive</u> constant c_0 , there exist n_0 s.t.: $c_0 g(n) < f(n)$ for all $n \geq n_0$	$n^2 + 100n = \omega(n\sqrt{n})$ $25n^2 + 100n \neq \omega(n^2)$

Properties

1. $f(n) = O(g(n)) \Rightarrow g(n) = \Omega(f(n))$
2. $f(n) = \Omega(g(n)) \Rightarrow g(n) = O(f(n))$
3. $f(n) = \Theta(g(n)) \Rightarrow g(n) = \Theta(f(n))$
4. If $f(n) = O(g(n))$ and $f(n) = \Omega(g(n)) \Rightarrow f(n) = \Theta(g(n))$
5. If $f(n) = \Theta(g(n)) \Rightarrow f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$

Transitivity (proved in slides):

6. If $f(n) = O(g(n))$ and $g(n) = O(h(n))$, then $f(n) = O(h(n))$.
7. If $f(n) = \Omega(g(n))$ and $g(n) = \Omega(h(n))$, then $f(n) = \Omega(h(n))$.

$1/n, 1, \lg n, n^\varepsilon, \sqrt{n}, n, n \lg n, n^2, n^3, n^d, c^n, n!, n^n$
where $0 < \varepsilon < 0.5$ and c and d are constants

Substitution method: If $\lim_{x \rightarrow \infty} h(x) = \infty$, and $h(x)$ is monotonically increasing
then $f(x) = O(g(x)) \Rightarrow f(h(x)) = O(g(h(x)))$. (topic not required)

Notation abuse:

Instead of $f(n) \in \Theta(g(n))$
we use: $f(n) = \Theta(g(n))$

$$a^{\log_b(n)} = n^{\log_b(a)} \text{ but } (a^n \neq n^a)$$


(E.g. $9^{\log_3 N} = N^{\log_3 9} = N^2$)

If $0 \leq c < d$, then $n^c = o(n^d)$. (E.g. $n^2 = o(n^5)$)

(Higher-order polynomials grow faster than lower-order ones.)

For any d , if $c > 1$, $n^d = o(c^n)$ (E.g. $n^{100} = o(2^n)$, b.c. $\lim_{n \rightarrow \infty} \frac{n^{100}}{2^n} = 0$)

(Exponential functions grow faster than polynomial ones.)



Typically, $f(n)$ is the running time of an algorithm. ($f(n)$ can be a complicated function.)

We try to find a $g(n)$ that is simple (e.g. n^2), and bounds $f(n)$. E.g. $f(n) = \Theta(g(n))$.